



Palliative Performance Scale (PPS)

| % | Ambulation | Activity Level and Evidence of Disease | Self-Care | Intake | Level of Consciousness |
|-----|------------------|---|--------------------------------|-------------------|-----------------------------|
| 100 | Full | <ul style="list-style-type: none"> • Normal • No disease evident | Full | Normal | Full |
| 90 | Full | <ul style="list-style-type: none"> • Normal • Some disease evident | Full | Normal | Full |
| 80 | Full | <ul style="list-style-type: none"> • Normal with effort • Some disease evident | Full | Normal or Reduced | Full |
| 70 | Reduced | <ul style="list-style-type: none"> • Unable to perform normal job/work • Some disease evident | Full | As above | Full |
| 60 | Reduced | <ul style="list-style-type: none"> • Unable to perform any job/work • Significant disease | Occasional assistance needed | As above | Full or confusion |
| 50 | Mainly Sit / Lie | <ul style="list-style-type: none"> • Unable to perform any job/work • Extensive disease | Considerable assistance needed | As above | Full or confusion |
| 40 | Mainly in Bed | <ul style="list-style-type: none"> • As above | Mainly assistance | As above | Full or drowsy or confusion |
| 30 | Bed Bound | <ul style="list-style-type: none"> • As above | Total care | Reduced | As above |
| 20 | Bed Bound | <ul style="list-style-type: none"> • As above | As above | Minimal | As above |
| 10 | Bed Bound | <ul style="list-style-type: none"> • As above | As above | Mouth care only | Drowsy or coma |
| 0 | Death | | | | |